

Principle of minimum complexity as a new principle in hadronic scattering

D. B. Ion and M. L. D. Ion

2 Dec 2005

Abstract

In this paper a measure of complexity of the system of $[J$ and $\theta]$ -quantum states produced in hadronic scattering is introduced in terms of the $[S_J(p), S_\theta(q)]$ -scattering entropies. We presented strong experimental evidence for the *saturation of the $[S_J^{o1}(p), S_\theta^{o1}(q)]$ -PMD-SQS optimal limits* in the pion-nucleon and kaon-nucleon scatterings. The validity of a *principle of minimum complexity in hadronic interaction* is supported with high accuracy ($CL > 99\%$) by the experimental data on pion-nucleon especially at energies higher than 2 GeV.

1 Introduction

In this paper, by using the optimality [1], the scattering entropies [2-4] $[S_J(p), S_\theta(q)]$, a measure of the complexity [5] of quantum scattering in terms of Tsallis-like entropies [6] is proposed. Then, the *nonextensive statistical behavior, optimality [1] and complexity of the $[J$ and $\theta]$ -quantum states* in hadronic scatterings are investigated in an unified manner. A connection between optimal states obtained from the *principle of minimum distance in the space of quantum states* (PMD-SQS)[1] and the most stringent (MaxEnt) entropic bounds on Tsallis-like entropies for quantum scattering is established. A measure of the complexity of quantum scattering in terms of Tsallis-like entropies is proposed. The results on the experimental tests of *the PMD-SQS-optimality*, as well as on the *complexity*, obtained by using the experimental pion-nucleon pion-nucleus phase shifts, are presented. Then, the nonextensivity indices p and q are determined from the experimental entropies by a

fit with the optimal entropies $[S_J^{ol}(p), S_\theta^{ol}(q)]$ obtained from the *principle of minimum distance in the space of states*. In this way strong experimental evidences for the p -nonextensivities in the range $0.5 \leq p \leq 0.6$ with q correlated via relation $1/p + 1/q = 2$, are obtained with high accuracy ($CL > 99\%$) from the experimental data of the pion-nucleon and pion-nucleus scatterings.

2 What is the complexity of a quantum system of scattering states?

In the last time many authors have developed and commented on the multiplicity and variety of definitions of *complexity* (C) in the scientific literature (see Ref.[5] and references therein).

What is the *complexity* of an interacting system? In general complexity is a concept very difficult to be defined. If we consider that this notion is coming from the original Latin word *complexus*, which signifies "*entwined*", "*twisted together*" then this may interpreted in the following way: in order to have a complex system you need two or more distinct components, which are joined in such a way that it is difficult to separate them. If we look in the Oxford Dictionary we find that *something is defined as complex if it is "made of closely connected parts"*. So, in these definitions we find the basic duality between parts which are at the same time distinct and connected. This implies that: a system would be more complex if more parts could be distinguished, and if more connections between them exist. So, the aspects of *distinction* and *connections* between the parts of systems determine two dimensions characterizing complexity. Distinction corresponds to the fact that different parts of the complex behaves differently while connections correspond to constrains. Moreover, the *distinction* leads at the limits to *disorder* (Δ), or chaos, while the *connection* leads to order (Ω). Complexity can only exist if both aspects are present: neither perfect disorder nor perfect order are complex. Before of a precise mathematical definition of the *complexity measure* (C) it is useful to point out some intuitive properties of the *complexity* of the physical systems [5]. These are as follows:

- (a) Any reasonable *measure of complexity must vanish for the complete ordered or complete disordered states of the system*;
- (b) Any *complexity must be an universal physical quantity*—that is, it is applied to any dynamical system;

(c) Any *complexity must correspond to mathematical complexity in problem solving.*

Current definitions may be divided into three distinct categories according to the following essential feature: (a) Definitions which take *complexity* as *monotonically increasing function* of *disorder* (Δ); (b) Complexity defined as a *convex function* of *disorder* (Δ); Complexity defined as *monotonically increasing function* of *order* ($\Omega = 1 - \Delta$).

Since we will express our *complexity measure* (C) in terms of *disorder* (Δ), or of *order* (Ω), we start with the definitions [5]

$$\Delta = \frac{S}{S_{\max}}, \quad \Omega = 1 - \Delta = 1 - \frac{S}{S_{\max}} \quad (1)$$

where S is the Boltzmann-Gibbs-Shannon (BGS) entropy

$$S = - \sum p_i \ln p_i \quad (2)$$

where $p_i, 1 \leq i \leq N$, is the probability of i -state of the N states available to the system (the Boltzmann constant k is taken $k=1$). In the simplest case of an isolated system, the maximum entropy occurs to the equiprobable distribution $p_i = 1/N, 1 \leq i \leq N$, yielding

$$S_{\max} = \ln N \quad (3)$$

All categories of *complexity measures* can be included by defining a measure of the form [5]

$$C^{\alpha\beta}(q) = \Delta^\alpha \Omega^\beta \quad (4)$$

So, the *complexity measures* of category I and III correspond to the particular cases: $\alpha > 0, \beta = 0$, and $\alpha = 0, \beta > 0$. The category II is obtained when both $\alpha > 0$ and $\beta > 0$. Then, the complexity measures $C^{\alpha\beta}(q)$ vanishes at zero *disorder* and zero *order*, and has the maximum

$$C_{\max}^{\alpha\beta} = \frac{\alpha^\alpha \beta^\beta}{(\alpha + \beta)^{(\alpha+\beta)}} \text{ at } \Delta_m = \frac{\alpha}{\alpha + \beta}, \quad \Omega_m = \frac{\beta}{\alpha + \beta} \quad (5)$$

3 Entropies for two-body scattering

Therefore, in order to introduce a measure of complexity of the hadronic system of quantum states produced in hadronic scatterings we must define

the *disorder* (Δ), or *order* (Ω) for the hadron-hadron scattering. These physical quantities can be defined just as in Eqs (1)-(2) by using the scattering Boltzman-Gibs-like entropy [2] or more general Tsallis-like scattering entropies introduced by us in Refs. [1,2]. For completeness we start our discussion with $f_{++}(x)$ and $f_{+-}(x)$, $x \in [-1, 1]$, as the scattering helicity amplitudes of *meson-nucleon scattering*

$$M(0^-) + N\left(\frac{1}{2}^+\right) \rightarrow M(0^-) + N\left(\frac{1}{2}^+\right) \quad (6)$$

where $x = \cos(\theta)$, θ being the c.m. scattering angle. The normalization of the helicity amplitudes $f_{++}(x)$ and $f_{+-}(x)$ is chosen such that the c.m. differential cross section $\frac{d\sigma}{d\Omega}(x)$ is given by

$$\frac{d\sigma}{d\Omega}(x) = |f_{++}(x)|^2 + |f_{+-}(x)|^2 \quad (7)$$

Since we will work at fixed energy, the dependence of σ_{el} and $\frac{d\sigma}{d\Omega}(x)$ and of $f(x)$, on this variable was suppressed. Hence, the helicities of incoming and outgoing nucleons are denoted by μ, μ' , and was written as $(+), (-)$, corresponding to $(\frac{1}{2})$ and $(-\frac{1}{2})$, respectively. In terms of the partial waves amplitudes f_{J+} and f_{J-} we have

$$\begin{aligned} f_{++}(x) &= \sum_{J=\frac{1}{2}}^{J_{\max}} (J+1/2)(f_{J-} + f_{J+})d_{\frac{1}{2}\frac{1}{2}}^J(x) \\ f_{+-}(x) &= \sum_{J=\frac{1}{2}}^{J_{\max}} (J+1/2)(f_{J-} - f_{J+})d_{-\frac{1}{2}\frac{1}{2}}^J(x) \end{aligned} \quad (8)$$

where the $d_{\mu\nu}^J(x)$ -rotation functions are given by

$$d_{\frac{1}{2}\frac{1}{2}}^J(x) = \frac{1}{l+1} \cdot \left[\frac{1+x}{2}\right]^{\frac{1}{2}} \left[\overset{\circ}{P}_{l+1}(x) - \overset{\circ}{P}_l(x)\right] \quad (9)$$

$$d_{-\frac{1}{2}\frac{1}{2}}^J(x) = \frac{1}{l+1} \cdot \left[\frac{1-x}{2}\right]^{\frac{1}{2}} \left[\overset{\circ}{P}_{l+1}(x) + \overset{\circ}{P}_l(x)\right] \quad (10)$$

$P_l(x)$ are Legendre Polynomials and prime indicates differentiation with respect to $x \equiv \cos \theta$.

Now, the elastic integrated cross section for the meson-nucleon scattering is expressed in terms of partial amplitudes f_{J+} and f_{J-}

$$\sigma_{el}/2\pi = \int_{-1}^{+1} dx \frac{d\sigma}{d\Omega}(x) = \sum_{J=\frac{1}{2}}^{J_{\max}} (2J+1)(|f_{J+}|^2 + |f_{J-}|^2) \quad (11)$$

J-nonextensive entropy for the quantum scattering

Now, we define two kind of Tsallis-like scattering entropies. One of them, namely $S_J(p)$, $p \in R$, is special dedicated to the investigation of the nonextensive statistical behavior of the *angular momentum J-quantum states*, and can be defined by

$$S_J(p) = \left[1 - \sum (2l+1)p_l^p \right] / (p-1), \quad p \in R, \quad (12)$$

where the probability distributions p_J , is given by

$$p_J = \frac{|f_{J+}|^2 + |f_{J-}|^2}{\sum_{J=\frac{1}{2}}^{J_{\max}} (2J+1)(|f_{J+}|^2 + |f_{J-}|^2)}, \quad \sum_{J=\frac{1}{2}}^{J_{\max}} (2J+1)p_J = 1 \quad (13)$$

θ -nonextensive entropy for the quantum scattering

In similar way, for the θ -scattering states considered as statistical canonical ensemble, we can investigate their nonextensive statistical behavior by using an angular Tsallis-like scattering entropy $S_\theta(q)$ defined as

$$S_\theta(q) = \left[1 - \int_{-1}^{+1} dx [P(x)]^q \right] / (q-1), \quad q \in R \quad (14)$$

where

$$P(x) = \frac{2\pi}{\sigma_{el}} \cdot \frac{d\sigma}{d\Omega}(x), \quad \int_{-1}^{+1} P(x)dx = 1 \quad (15)$$

with $\frac{d\sigma}{d\Omega}(x)$ and σ_{el} defined by Eqs.(7)-(11).

The above Tsallis-like scattering entropies posses two important properties. First, in the limit $k \rightarrow 1$, $k \equiv p, q$, the Boltzman-Gibbs kind of entropies is recovered:

$$\begin{aligned} \lim_{p \rightarrow 1} S_J(p) &= S_J(1) = -\sum (2J+1)p_J \ln p_J \\ \lim_{q \rightarrow 1} S_\theta(q) &= S_\theta(1) = -\int_{-1}^{+1} dx P(x) \ln P(x) \end{aligned} \quad (16)$$

Secondly, these entropies are *nonextensive* in the sense that

$$S_{A+B}(k) = S_A(k) + S_B(k) + (1-k)S_A(k)S_B(k), \quad k = p, q \quad (17)$$

for any independent sub-systems A, B ($p_{A+B} = p_A \cdot p_B$). Hence, each of the indices $p \neq 1$ or $q \neq 1$ from the definitions (12) and (17) can be interpreted as measuring the degree *nonextensivity of the quantum state system*.

We next consider the *maximum-entropy* (MaxEnt) problem

$$\max\{S_J(p), S_\theta(q)\} \text{ when } \sigma_{el} = \text{fixed and } \frac{d\sigma}{d\Omega}(1) = \text{fixed} \quad (18)$$

as criterion for the determination of the *equilibrium distributions* p_l^{me} and $P^{me}(x)$ for the quantum states from the $(0^{-\frac{1}{2}} \rightarrow 0^{-\frac{1}{2}})$ -scattering. For the J -*quantum states*, in the spin $(0^{-\frac{1}{2}} \rightarrow 0^{-\frac{1}{2}})$ scattering case, these distributions are given by:

$$\begin{aligned} p_J^{me} &= p_J^{o1} = \frac{1}{2K_{\frac{1}{2}\frac{1}{2}}(1,1)} = \frac{1}{(J_o+1)^2-1/4}, \text{ for } \frac{1}{2} \leq J \leq J_o, \text{ and} \\ p_J^{me} &= 0, \text{ for } J \geq J_o + 1 \end{aligned} \quad (19)$$

while, for the θ -*quantum states*, these distributions are as follows

$$P^{me}(x) = P^{o1}(x) = \left[\frac{K_{\frac{1}{2}\frac{1}{2}}^2(x, 1)}{K_{\frac{1}{2}\frac{1}{2}}(1, 1)} \right] \quad (20)$$

Consequently, by a straightforward calculus we obtain that the solution of the problem (18) is given by

$$S_J^{\max}(p) = S_J^{o1}(p) = \left[1 - [2K_{\frac{1}{2}\frac{1}{2}}(1, 1)]^{1-p} \right] / (p - 1), \text{ for } p > 0 \quad (21)$$

$$S_\theta^{\max}(q) = S_\theta^{o1}(q) = \left[1 - \int_{-1}^{+1} dx \left(\frac{[K_{\frac{1}{2}\frac{1}{2}}(x, 1)]^2}{K_{\frac{1}{2}\frac{1}{2}}(1, 1)} \right)^q \right] / (q - 1), \text{ for } q > 0 \quad (22)$$

where the *reproducing kernel function* $K_{\frac{1}{2}\frac{1}{2}}(x, 1)$ is given by

$$K_{\frac{1}{2}\frac{1}{2}}(x, 1) = \frac{1}{2} \sum_{1/2}^{J_o} (2J + 1) d_{\frac{1}{2}\frac{1}{2}}^J(x) \quad (23)$$

while the *optimal angular momentum* J_o is

$$(J_o + 1)^2 - 1/4 = 2K_{\frac{1}{2}\frac{1}{2}}(1, 1) = \frac{4\pi}{\sigma_{el}} \frac{d\sigma}{d\Omega}(1) \quad (24)$$

$d_{\frac{1}{2}\frac{1}{2}}^J(x)$ are the usual spin 1/2-rotation functions.

Proof: The proof of the problem (18) is equivalent to the following unconstrained maximization problem:

$$\mathcal{L} \rightarrow \max \quad (25)$$

where the Lagrangian function is defined as

$$\begin{aligned} \mathcal{L} \equiv & \lambda_0 \{S_J(p), S_\theta(q)\} + \lambda_1 \left\{ \sigma_{el}/4\pi - \sum (2J+1) \left[|f_{J-}|^2 + |f_{J+}|^2 \right] \right\} \\ & \lambda_2 \left\{ \frac{d\sigma}{d\Omega}(1) - \left[\sum (2J+1) \operatorname{Re} f_{J+} \right]^2 - \left[\sum (2J+1) \operatorname{Im} f_{J+} \right]^2 \right\} \end{aligned} \quad (26)$$

Hence, the solution of the problem (18), correspond to the the singular case $\lambda_0 = 0$, and is reduced just to the solution of the *minimum constrained distance in space of quantum states*:

$$\min \sum (2J+1) \left[|f_{J-}|^2 + |f_{J+}|^2 \right] \text{ when } \frac{d\sigma}{d\Omega}(1) = \text{is fixed} \quad (27)$$

from which we obtain the following analytic form of the optimal scattering helicity amplitudes

$$f_{++}^{o1}(x) = f_{++}(1) \cdot \frac{K_{\frac{1}{2}\frac{1}{2}}(x, 1)}{K_{\frac{1}{2}\frac{1}{2}}(1, 1)}, \quad f_{+-}^{o1}(x) = 0, \quad (28)$$

where the *reproducing kernels* $K_{\frac{1}{2}\frac{1}{2}}(x, 1)$ are given by Eqs. (23)-(24). All these results together other are presented in Table 1

4 Complexity measure for quantum scattering

Now, let us apply the definitions (1) and (4) of the *disorder*, *order*, and *complexity measure* to the nonextensive quantum scattering of the elementary particles. Then, we define the scattering (J, θ) -*disorders* as follows

$$\Delta_X(q) = \frac{S_X(q)}{S_X^{\max}(q)}, \quad X \equiv \theta, J \quad (29)$$

and the corresponding scattering (θ, J) -orders by the relations

$$\Omega_X(q) = 1 - \Delta_X(q), \quad X \equiv \theta, J \quad (30)$$

where $S_\theta(q), S_J(q)$ are the scattering *Tsallis-like scattering entropies* defined in Section 3.

Therefore, using Eqs. (29)-(30) and (12)-(15) we can calculate the *scattering complexities* $C_X^{\alpha\beta}(q)$ defined as follows

$$C_X^{\alpha\beta}(q) = \Delta_X^\alpha(q) \Omega_X^\beta(q), \quad X \equiv \theta, J \text{ and } q \in R \quad (31)$$

5 Numerical Results

Now, for a systematic experimental investigation of the saturation of the *optimality limits in hadron-hadron scattering* is necessary to use the formulas from the Table 1 and the available experimental phase-shifts [8-10] to solve the following numerical problems: (i) To reconstruct the experimental pion-nucleon, kaon-nucleon and antikaon-nucleon scattering amplitudes; (ii) To obtain numerical values of the experimental scattering entropies $S_J(q), S_\theta(q), \bar{S}_{J\theta}(p, q)$ from the reconstructed amplitudes; (iii) To obtain the numerical values of the *optimal angular momentum* J_o from experimental scattering amplitudes and to calculate the numerical values for the PMD-SQS-optimal entropies $S_J^{o1}(q), S_\theta^{o1}(q); \bar{S}_{J\theta}(p, q)$; (iv) To obtain numerical values for $\chi_J^2(p)$ or/and $\chi_\theta^2(q)$ -test functions given by

$$\chi_X^2(k) = \sum_{i=1}^{n_{\text{exp}}} \left[\frac{[S_X(k)]_i - [S_X^{o1}(k)]_i}{[\Delta S_X^{o1}]_i} \right]^2, \quad X \equiv J, \theta; \bar{J}\theta \quad k \equiv p, q \quad (32)$$

where

$$\Delta S_X^{o1}(k) = |[S_X^{o1}(k)]_{J_o+1} - [S_X^{o1}(k)]_{J_o-1}| \quad (33)$$

are the values of the *PMD-SQS-optimal entropies* $[S_X^{o1}(k)]_{J_o \pm 1}$ calculated with the optimal angular momenta $J_o \pm 1$, respectively. This procedure is equivalent with assumption of an error of $\Delta J_o = \pm 1$ in estimation of the experimental values of the optimal angular momentum J_o . The results obtained in this way are presented in the Figs. 1 and Tables 2-4. So, the best fit is obtained (see Tables 2) for the correlated pairs p and $q = p/(2p - 1)$ with the values of p in the range $p = 0.6$ and $q = p/(2p - 1) = 3.0$. However, at

this level of error analysis the case of the extensive system ($p=q=1$) of quantum scattering states is also consistent with the experimental data while the pair of nonextensivities ($p=3.0, q=0.6$) rejected by the values of χ^2/n_D -test function (see Table 4).

Moreover, in Fig. 2-3 we presented the numerical results for the *complexity of the J-quantum systems of states* produced in the pion-nucleon scattering. So, the experimental values of the complexities $C_J^{11}(p)$ for π^+p and π^-p -scatterings, obtained from the available experimental phase-shifts [19], are compared with the *PMD-SQS-optimal state* predictions (full curve $[C_J^{11}(p)]^{o1} = 0$). In all cases, the error-bands (the grey regions) around the optimality curves are calculated by assuming an error of $\Delta J_o = \pm 1$ in calculation of the optimal angular momentum J_o . From the results of the Figs.2-3 we conclude that the scattering complexities $C_J^{11}(p)$ are in excellent agreement with the optimal complexities $[C_J^{11}(p)]^{o1} = 0$. So, a *principle of minimum complexity in hadronic interaction* is suggested with high accuracy ($CL > 99\%$) by the experimental pion-nucleon scattering data.

6 Summary and Conclusions

In this paper, by using *the optimality* [1-4] and $[S_J(p), S_\theta(q)]$ -Tsallis-like entropies, the *complexity* and nonextensivity of the $[J$ and $\theta]$ -quantum states produced in hadronic scatterings are investigated. The main results and conclusions can be summarized as follows:

- (i) Using the available experimental phase shifts analysis we calculated the numerical values for the $[S_J(p), S_\theta(q)]$ -Tsallis-like scattering entropies for the pure isospin I -scattering states: $[(\pi N)_{I=1/2,3/2}; (KN)_{I=0,1}; (\bar{K}N)_{I=0,1}]$;
- (ii) We presented strong experimental evidence for the *saturation of the* $[S_J^{o1}(p), S_\theta^{o1}(q)]$ -PMD-SQS optimal limits for all nonextensive (J, θ) -statistical ensembles of quantum states produced in hadron-hadron scattering (see Fig.1 and Table 2). These results allow to conclude that the $[J]$ -quantum system and $[\theta]$ -quantum system are produced at "equilibrium" but with the $[\frac{1}{2p} + \frac{1}{2q}]$ -conjugated nonextensivities $p = 0.6$ and $q = p/(2p - 1) = 3$ in all investigated isospin scattering states: $[(\pi N)_{I=1/2,3/2}; (KN)_{I=0,1}; (\bar{K}N)_{I=0,1}]$. So the "geometric origin" of the nonextensivities p and q (as dimensions of the Hilbert spaces L_{2p} and

L_{2q}) as well as their correlations are experimentally confirmed with high accuracy ($CL > 99\%$). The strong experimental evidence obtained here for the *nonextensive statistical behavior* of the (J, θ) - quantum scatterings states in the pion-nucleon, kaon-nucleon and antikaon-nucleon scatterings can be interpreted as an indirect manifestation of *the presence of the quarks and gluons* as fundamental constituents of the scattering system having the strong-coupling long-range regime required by the Quantum Chromodynamics.

- (iii) We presented strong experimental evidence for the validity of *a principle of minimum complexity (PMC) in pion-nucleon scattering*. *PMC* is supported with high accuracy ($CL > 99\%$) by all the experimental data especially at energies higher than 2 GeV beyond the resonance region.

Finally, we note that further investigations are needed since this *saturation of optimality and complexity limits* as well as *nonextensive statistical behavior* of the quantum scattering, evidenced here with high accuracy ($CL > 99\%$), can be a signature for a *new universal law* of the quantum scattering.

References

- [1] D. B. Ion, Phys. Lett. **B 376**, 282 (1996).
- [2] D. B. Ion and M. L. D. Ion, Phys Rev. Lett. **81** (1998) 5714; M. L. D. Ion and D. B. Ion, Phys Rev. Lett. **83** (1999) 463.
D.B. Ion and M.L.D. Ion, Phys. Rev. **E 60**, 5261(1999); M.L.D. Ion and D.B. Ion, Phys. Lett. **B 482**,57 (2000); D.B. Ion and M.L.D. Ion, Phys. Lett. **B 503**, 263 (2001); D.B. Ion and M.L.D. Ion, Phys. Lett. **B 519**, 63 (2001); D.B. Ion and M.L.D. Ion, in *Classical and Quantum Complexity and Nonexten-sive Thermodynamics*, eds. P. Grigolini, C. Tsallis and B.J. West, Chaos , Solitons and Fractals **13** (2002) 547; D. B. Ion and M. L. Ion, Physica **A340** (2004) 501.
- [3] M.L.D. Ion and D.B. Ion, Phys. Lett. **B 474**, 395 (2000).

- [4] D. B. Ion, International J. Theor. Phys. **24**, 1217 (1985); D. B. Ion, International J. Theor. Phys. **25**, 1257 (1986); D. B. Ion, Rev. Rom. Phys. **26**, 15 and 25 (1981); D. B. Ion, Rev. Rom. Phys. **36**, 251 (1991).
- [5] J. S. Shiner, M. Davidson and Landsberg, Phys. Rev. E **59** (1999) 1459.
- [6] C. Tsallis, J. Stat. Phys. **52**, 479 (1988).
- [7] C. Tsallis and E. P. Borges, *Nonextensive statistical mechanics-Applications to nuclear and high energy physics*, **ArXiv**: cond-mat/0301521 (2003), to appear in the Proceedings of the *X International Workshop on Multiparticle Production Correlations and Fluctuations in QCD* (8-15 June 2002, Crete), ed. N. Antoniou (World Scientific, 2003); An updated bibliography on Nonextensive Statistics can be found at the website: <http://tsallis.cat.cbpf.br/biblio.htm>. For more general nonextensive entropies see: E. P. Borges and I. Roditi in Phys. Lett. **A246**, 399 (1998).
- [8] G. Höhler, F. Kaiser, R. Koch, and E. Pietarinen, *Physics Data, Handbook of Pion-Nucleon Sattering*, 1979, **Nr.12-1**
- [9] R. A. Arndt and L. D. Roper, Phys. Rev. **D31**, 2230 (1985).
- [10] M. Alston-Garnjost, R.W. Kenney, D.L. Pollard, R.R. Ross, R.D. Tripp, H. Nicholson, M. Ferro-Luzzi, Phys. Rev. **D18**, 182 (1978)
- [11] L. R. Lopez, L.J. Caufield, *A Principle of Minimum Complexity in Evolution*, Lecture Notes in Computer Science, **496**, 405 (1991).

TABLE CAPTIONS

- Table 1:** The optimal distributions, reproducing kernels, optimal entropies, entropic bands, for the $(0^-1/2^+ \rightarrow 0^-1/2^+)$ -scatterings.
- Table 2:** χ^2/n_D obtained from comparisons of the experimental scattering entropies: $S_J(p)$, $S_\theta(q)$ with the optimal entropies: $S_J^{o1}(p)$, $S_\theta^{o1}(q)$, respectively, for the pair $(p=0.6, q=3.0)$.
- Table 3:** χ^2/n_D obtained from comparisons of the experimental scattering entropies: $S_J(p)$, $S_\theta(q)$ with the optimal entropies: $S_J^{o1}(p)$, $S_\theta^{o1}(q)$, respectively, for the pair $(p=1.0, q=1.0)$.
- Table 4:** χ^2/n_D obtained from comparisons of the experimental scattering entropies: $S_J(p)$, $S_\theta(q)$ with the optimal entropies: $S_J^{o1}(p)$, $S_\theta^{o1}(q)$, respectively, for the pair $(p=3.0, q=0.6)$.

FIGURES CAPTIONS

- Fig. 1:** The experimental values of the Tsallis-like scattering entropies $S_J(p)$ (12) for $[(\pi N)_{I=1/2,3/2}; (KN)_{I=0,1}; (\overline{K}N)_{I=0,1}]$ -scatterings, obtained from the available experimental phase-shifts [8-10], are compared with the PMD-SQS-optimal state predictions $S_J^{o1}(p)$ (full curve). The error-band (the grey region) around the optimality curve is calculated by assuming an error of $\Delta J_o = \pm 1$ in calculation of the optimal angular momentum J_o .
- Fig. 2:** The experimental values of the complexities $C_J^{11}(p)$ for π^+p -scatterings, obtained, by using Eqs. (29)-(31), from the available experimental phase-shifts [8-10], are compared with the PMD-SQS-optimal state complexity (full curve $C_J^{11}(p) = 0$). The error-band (the grey region) around the optimality curve is calculated by assuming an error of $\Delta J = \pm 1$ in calculation of the optimal angular momentum J_o .
- Fig. 3:** The experimental values of the complexities $C_J^{11}(p)$ for π^-p -scatterings, obtained from the available experimental phase-shifts [8], are compared with the PMD-SQS-optimal state complexity (the full curve $C_J^{11}(p) = 0$). The error-band (the grey region) around the optimality curve is calculated by assuming an error of $\Delta J = \pm 1$ in calculation of the optimal angular momentum J_o .

Table 1

Nr.	Name	$(0^-1/2^+ \rightarrow 0^-1/2^+) - \text{scatterings}$
1	Optimal inequalities	$\frac{d\sigma}{d\Omega}(x) \leq K_{\frac{1}{2}\pm\frac{1}{2}}(\pm 1, \pm 1) \ f\ ^2$
2	Optimal states	$f^{o+1}(x) = f(1) \frac{K_{\frac{1}{2}\frac{1}{2}}(x,+1)}{K_{\frac{1}{2}\frac{1}{2}}(1,1)}, f^{o+1}(x) = f(-1) \frac{K_{-\frac{1}{2}\frac{1}{2}}(x,-1)}{K_{-\frac{1}{2}\frac{1}{2}}(-1,-1)}$
2	Reproducing kernels $K_{\frac{1}{2}\pm\frac{1}{2}}(x, \pm 1)$	$K_{\frac{1}{2}\frac{1}{2}}(x, +1) = \sum_{\frac{1}{2}}^{J_o} (J + \frac{1}{2}) d_{\frac{1}{2}\frac{1}{2}}^J(x)$ $K_{\frac{1}{2}-\frac{1}{2}}(x, -1) = \sum_{\frac{1}{2}}^{J_o} (J + \frac{1}{2}) d_{\frac{1}{2}-\frac{1}{2}}^J(x)$ $2K_{\frac{1}{2}\pm\frac{1}{2}}(\pm 1, \pm 1) = (J_o + 1)^2 - 1/4$
3	Optimal distributions $P^{o\pm 1}(x)$	$P^{o1}(x) = \frac{[K_{\frac{1}{2}\frac{1}{2}}(x,1)]^2}{K_{\frac{1}{2}\frac{1}{2}}(1,1)}, P^{o-1}(x) = \frac{[K_{\frac{1}{2}-\frac{1}{2}}(x,-1)]^2}{K_{\frac{1}{2}-\frac{1}{2}}(-1,-1)}$
4	Optimal distribution $\{p_J^{o1}\}$	$p_J^{o\pm 1} = \frac{1}{2K_{\frac{1}{2}\pm\frac{1}{2}}(\pm 1, \pm 1)}, \text{ for } 1/2 \leq J \leq J_o$ $p_J^{o\pm 1} = 0 \text{ for } J \geq J_o + 1$
5	Number of optimal states for $y = 1$	$N_o = \sum (2J + 1) = (J_o + 1)^2 - 1/4 = 2K_{\frac{1}{2}\pm\frac{1}{2}}(\pm 1, \pm 1)$
6	Optimal angular – momentum	$J_o = \left\{ \left[\frac{4\pi}{\sigma_{el}} \frac{d\sigma}{d\Omega}(\pm 1) + \frac{1}{4} \right]^{1/2} - 1 \right\}$
7	Optimal entropy $S_J^{o1}(q)$	$S_J^{o\pm 1}(p) = \frac{1}{p-1} [1 - N_{os}^{1-p}]$
8	Optimal entropy $S_\theta^{o1}(q)$	$S_\theta^{o\pm 1}(q) = \frac{1}{q-1} \left[1 - \int_{-1}^{+1} dx \left(\frac{[K_{\frac{1}{2}\pm\frac{1}{2}}(x, \pm 1)]^2}{K(\pm 1, \pm 1)} \right)^q \right]$
11	J –entropic band	$0 \leq S_J(p) \leq S_J^{o\pm 1}(p)$
12	θ –entropic band	$\frac{1}{q-1} [1 - K(1, 1)^{q-1}] \leq S_\theta(q) \leq S_\theta^{o\pm 1}(q)$

Table 2

	Hadron-hadron scattering	Number of PSA	p	q= $\frac{p}{2p-1}$	χ^2/n_D $S_J(p)$ $S_\theta(q)$
1	$\pi N \rightarrow (\pi N)_{I=1/2}$	88	0.6	3.0	0.102 0.015
2	$\pi N \rightarrow (\pi N)_{I=3/2}$	88	0.6	3.0	0.130 0.090
3	$KN \rightarrow (KN)_{I=0}$	52	0.6	3.0	0.449 0.035
4	$KN \rightarrow (KN)_{I=1}$	53	0.6	3.0	0.089 0.045
5	$\bar{K}N \rightarrow (\bar{K}N)_{I=0}$	50	0.6	3.0	0.168 0.009
6	$\bar{K}N \rightarrow (\bar{K}N)_{I=1}$	50	0.6	3.0	0.062 0.010

Table 3

Hadron-hadron scattering	Number of PSA	p q= $\frac{p}{2p-1}$	χ^2/n_D S _J (p) S _θ (q)
$\pi N \rightarrow (\pi N)_{I=1/2}$	88	1 1	0.649 0.648
$\pi N \rightarrow (\pi N)_{I=3/2}$	88	1 1	0.691 1.059
$KN \rightarrow (KN)_{I=0}$	52	1 1	0.146 0.494
$KN \rightarrow (KN)_{I=1}$	53	1 1	0.259 0.586
$\bar{K}N \rightarrow (\bar{K}N)_{I=0}$	50	1 1	0.199 0.267
$\bar{K}N \rightarrow (\bar{K}N)_{I=1}$	50	1 1	0.064 0.196

Table 4

Hadron-hadron scattering	Number of PSA	p q= $\frac{p}{2p-1}$	χ^2/n_D S _J (p) S _θ (q)
$\pi N \rightarrow (\pi N)_{I=1/2}$	88	3.0 0.6	143.6 2.181
$\pi N \rightarrow (\pi N)_{I=3/2}$	88	3.0 0.6	209.2 3.010
$KN \rightarrow (KN)_{I=0}$	52	3.0 0.6	0.190 1.089
$KN \rightarrow (KN)_{I=1}$	53	3.0 0.6	30.53 1.030
$\bar{K}N \rightarrow (\bar{K}N)_{I=0}$	50	3.0 0.6	11.38 0.551
$\bar{K}N \rightarrow (\bar{K}N)_{I=1}$	50	3.0 0.6	16.00 0.391




Figure 1: The experimental values of the Tsallis-like scattering entropies $S_J(p)$ (12) for $[(\pi N)_{l=1/2,3/2}; (KN)_{l=0,1}; (\overline{KN})_{l=0,1}]$ –scatterings, obtained from the available experimental phase-shifts [8-10], are compared with the PMD-SQS-optimal state predictions $S_J^{o1}(p)$ (full curve). The error-band (the grey region) around the optimality curve is calculated by assuming an error of $\Delta J_o = \pm 1$ in calculation of the optimal angular momentum J_o .




Figure 2: The experimental values of the complexities $C_J^{11}(p)$ for π^+p -scatterings, obtained, by using Eqs. (29)-(31), from the available experimental phase-shifts [8-10], are compared with the PMD-SQS-optimal state complexity (full curve $C_J^{11}(p) = 0$). The error-band (the grey region) around the optimality curve is calculated by assuming an error of $\Delta J = \pm 1$ in calculation of the optimal angular momentum J_O .




Figure 3: The experimental values of the complexities $C_J^{11}(p)$ for π^-p -scatterings, obtained from the available experimental phase-shifts [8], are compared with the PMD-SQS-optimal state complexity (the full curve $C_J^{11}(p) = 0$). The error-band (the grey region) around the optimality curve is calculated by assuming an error of $\Delta J = \pm 1$ in calculation of the optimal angular momentum J_O .